

# Meet Kirk

54, police officer, husband, and father

Kirk struggled to manage his IgAN for over 5 years with medications and diet changes.\* When TARPEYO received approval, Kirk and his nephrologist discussed TARPEYO as an option.

\*TARPEYO is taken along with blood pressure medicine.

“ My nephrologist and I decided on 9 months of TARPEYO, in addition to my blood pressure medicine, to help treat my IgAN. My doctor was extremely satisfied with the outcome, and so was I. I’ve been off of the medication now for several months and my labs are still within range. ”

Kirk, *actual* TARPEYO patient



Scan this QR code to hear Kirk's story.

[tarpeyo.com/tarpeyo-testimonials](https://tarpeyo.com/tarpeyo-testimonials)



Kirk was compensated for his time.

## Important Safety Information (cont'd)

### How should I take TARPEYO?

- Take TARPEYO exactly as your healthcare provider tells you
- Your healthcare provider will decide how long you should take TARPEYO. Do not stop taking TARPEYO without first talking with your healthcare provider
- Take your prescribed dose of TARPEYO 1 time each day in the morning, at least 1 hour before a meal
- Swallow TARPEYO capsules whole. **Do not** open, chew, crush, or break TARPEYO capsules before swallowing
- If you miss a dose of TARPEYO, take your prescribed dose at your next scheduled time. **Do not** take two doses of TARPEYO at the same time
- If you take too much TARPEYO, call your healthcare provider right away or go to the nearest hospital emergency room

## Support and financial assistance made easy

### Support at every point of your journey with IgAN

**Touchpoints**  
Support beyond the script

TARPEYO Touchpoints is a program designed to support you on your treatment journey, offering services, assistance, and resources to help you access TARPEYO.

TARPEYO Touchpoints is available Monday through Friday, 8 AM to 8 PM ET, at 1-833-444-8277.

**97%** OF TARPEYO PATIENTS HAVE PAID LESS THAN \$10 PER PRESCRIPTION\*

\*All patients regardless of insurance type. This includes Commercial, Medicare, Medicaid, and cash-paying patients from January 2022 through November 2023.



### Stay connected with others in the IgAN community

The following advocacy groups offer support and valuable information about this rare disease.



**TAKE CHARGE. ASK YOUR DOCTOR IF TARPEYO IS RIGHT FOR YOU.**

## Important Safety Information (cont'd)

### What should I avoid while taking TARPEYO?

**Do not** eat grapefruit or drink grapefruit juice during your treatment with TARPEYO. Eating grapefruit or drinking grapefruit juice can increase the level of TARPEYO in your blood.

**Please see Important Safety Information throughout and accompanying Full Prescribing Information.**

**TARPEYO**  
(budesonide) delayed release capsules - 4 mg

## Important Safety Information (cont'd)

### What are the possible side effects of TARPEYO?

TARPEYO may cause serious side effects, including:

- **Effects of having too much corticosteroid medicine in your blood (hypercorticism):** Long-time use of TARPEYO can cause you to have signs and symptoms of too much cortisol, a stress hormone in your blood. Tell your healthcare provider if you have any of the following signs and symptoms of hypercorticism: acne, bruise easily, rounding of your face (moon face), ankle swelling, thicker or more hair on your body and face, a fatty pad or hump between your shoulders (buffalo hump), or pink or purple stretch marks on the skin of your abdomen, thighs, breasts, or arms
- **Adrenal suppression:** When TARPEYO is taken for a long period of time (chronic use), adrenal suppression can happen. This is a condition in which the adrenal glands do not make enough steroid hormones. Symptoms of adrenal suppression include tiredness, weakness, nausea and vomiting, and low blood pressure. Tell your healthcare provider if you are under stress or have any symptoms of adrenal suppression during treatment with TARPEYO
- **Risk of immunosuppression:** TARPEYO weakens your immune system. Taking medicines that weaken your immune system makes you more likely to get infections. Avoid contact with people who have contagious diseases, such as chickenpox or measles, during treatment with TARPEYO. Tell your healthcare provider right away if you come in contact with anyone who has chickenpox or measles. Consult with your healthcare provider regarding appropriate vaccination scheduling
- Tell your healthcare provider if you develop any symptoms of infection during treatment with TARPEYO, including fever, feeling tired, chills, aches, pain, and nausea and vomiting

### The most common side effects of TARPEYO include:

- swelling of the lower legs, ankles, and feet
- high blood pressure
- muscle spasms
- acne
- headache
- upper respiratory tract infection
- swelling of the face
- weight increase
- indigestion
- irritation or inflammation of the skin
- joint pain
- increased white blood cell count

These are not all the possible side effects of TARPEYO. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**Please see Important Safety Information throughout and accompanying Full Prescribing Information.**

**calliditas**  
THERAPEUTICS

TARPEYO is a registered trademark of Calliditas Therapeutics AB, or its affiliates. © 2024 Calliditas Therapeutics AB. All rights reserved. 03/24 US-TAR-2400006

**TARPEYO**  
(budesonide) delayed release capsules - 4 mg



Bill, Cathrin, and Kirk are *actual* TARPEYO patients.

Bill, Cathrin, and Kirk were compensated for their time.

## REAL PEOPLE AND RESILIENT STORIES OF PATIENTS WHO TOOK CHARGE WITH TARPEYO

TARPEYO is the first and only FDA-approved treatment to reduce the loss of kidney function in adults with IgA Nephropathy (IgAN)

## Indication

### What is TARPEYO?

TARPEYO is a prescription medicine used to reduce the loss of kidney function in adults with a kidney disease called primary immunoglobulin A nephropathy (IgAN) who are at risk for their disease getting worse. It is not known if TARPEYO is safe and effective in children.

## Important Safety Information

**Do not take TARPEYO** if you are allergic to budesonide or any of the ingredients in TARPEYO. See the end of the Patient Information for a complete list of ingredients in TARPEYO.

**Please see Important Safety Information throughout and accompanying Full Prescribing Information.**



# Meet Cathrin

37, astrophotographer and proud dog mom

After her diagnosis in 2011, Cathrin struggled to find a treatment that helped control her IgAN. She and her husband plan to expand their family, making her health and proactive management of IgAN more important than ever. That is why she chose to talk to her doctor about a 9-month course of TARPEYO.\*

\*TARPEYO is taken along with blood pressure medicine.



“My doctor suggested TARPEYO, and I was really impressed by the data. Specifically, the results from people who had finished their 9-months of treatment. Knowing that my kidney function could stay stable for some time made a difference for me.”

Cathrin, *actual* TARPEYO patient

See clinical trial results at [Tarpeyo.com](https://tarpeyo.com)



Scan this QR code to hear Cathrin's story.

[tarpeyo.com/tarpeyo-testimonials](https://tarpeyo.com/tarpeyo-testimonials)

Cathrin was compensated for her time.

## Important Safety Information (cont'd)

**Before taking TARPEYO, tell your healthcare provider about all of your medical conditions, including if you:**

- have liver problems
- plan to have surgery
- have chickenpox or measles or have recently been near anyone with chickenpox or measles
- have an infection
- have high blood sugar levels (prediabetes or diabetes)
- have glaucoma or cataracts
- have a family history of diabetes or glaucoma
- have or have had tuberculosis
- have high blood pressure (hypertension)
- have decreased bone mineral density (osteoporosis)

**Please see Important Safety Information throughout and accompanying Full Prescribing Information.**

# Meet Bill

52, preschool owner, father, and grandfather

Shortly after his IgAN diagnosis, Bill's disease began to progress rapidly. After his lab results came in, his doctor told him about TARPEYO.\* He was glad to hear that there was a disease-specific treatment option that may help him get ahead of his disease progression.†

\*TARPEYO is taken along with blood pressure medicine.

†It has not been established to what extent the efficacy of TARPEYO is from local effects (in the gut) vs systemic effects (in circulation).

“Getting ahead of my disease by targeting a source of IgAN was important to me in a treatment. I didn't have to watch and wait for my disease to progress.”

Bill, *actual* TARPEYO patient



Bill was compensated for his time.



Scan this QR code to hear Bill's story.

[tarpeyo.com/tarpeyo-testimonials](https://tarpeyo.com/tarpeyo-testimonials)

## Important Safety Information (cont'd)

**Before taking TARPEYO, tell your healthcare provider about all of your medical conditions, including if you:**

- have stomach ulcers
- are pregnant or plan to become pregnant. TARPEYO may harm your unborn baby. Talk to your healthcare provider about the possible risk to your unborn baby if you take TARPEYO when you are pregnant
- are breastfeeding or plan to breastfeed. It is not known if TARPEYO passes into your breast milk or if it will affect your baby. Talk to your healthcare provider about the best way to feed your baby during treatment with TARPEYO

**Tell your healthcare provider about all the medicines you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements. TARPEYO and other medicines may affect each other causing side effects.